

Table 1.1. Adult return goal to each subbasin. Most goals represent a combination of spawning needs plus tributary harvest opportunities; exceptions are footnoted.

SUBBASIN	CHINOOK			COHO	SOCK-EYE	STEELHEAD		TOTAL
	Spring	Summer	Fall			Summer	Winter	
<i>LOWER COLUMBIA RIVER MAINSTEM (Bonneville Dam → McNary Dam)</i>								
Wind River ^a	5,000			200		2,000	200	7,400
Little White River ^a	2,000		200	200				2,400
Big White River ^a	500		100	100		4,800	800	6,300
Klickitat River ^b	20,000		40,000	50,000		25,000		135,000
Hood River ^b	1,700					8,000	5,000	14,700
Deschutes River ^b	10,250		11,000		5,000	19,000		45,250
Fifteenmile Creek ^c							300	300
John Day River ^b	7,000					45,000		52,000
<i>MID-COLUMBIA RIVER MAINSTEM (McNary Dam → Chief Joseph Dam)</i>								
Umatilla River ^b	11,000		21,000	6,000		9,670		47,670
Hanford Reach ^c			40,000					40,000
Walla Walla River ^b	5,000					11,000		16,000
Yakima River ^b	26,300	12,000	4,700	5,000		29,700		77,700
Wenatchee River ^b	21,000	10,000			35,000	6,410		72,410
Entiat River ^a	750					3,000		3,750
Methow River ^a	2,000	3,000				10,000		15,000
Okanogan River ^a	1,000	2,000			15,000	10,000		28,000
<i>LOWER-SNAKE RIVER MAINSTEM (Ice Harbor Dam → Hells Canyon Dam)</i>								
Snake River ^b			18,300					18,300
Tucannon River ^b	3,000	2,000				2,200		7,200
Clearwater River ^b	60,000	48,000	6,000	4,500		93,100		211,600
Grande Ronde River ^b	16,000		10,000	3,500	2,500	27,500		59,500
Salmon River								NA
Imnaha River ^b	5,740		1,150			4,315		11,205
TOTAL	198,240	77,000	152,450	69,500	57,500	310,695	6,300	871,685

- ^a Harvest goal only
- ^b Harvest and spawning goal
- ^c Spawning goal only