

Table 1.2. Survival rates of Snake River spring and fall chinook under NMFS and Tribal restoration plans. *From Table 5C.2.*

| LIFE HISTORY STAGE OR MORTALITY SOURCE | SPRING CHINOOK | | FALL CHINOOK | |
|--|-------------------|--------------------------------------|-------------------|--------------------------------------|
| | NMFS 1995 Plan | Tribal 25-yr Strategy Option 3 | NMFS 1995 Plan | Tribal 25-yr Strategy Option 3 |
| Egg→Smolt | .045 | .10 | .50 | .50 |
| Juvenile Passage | .24 | .71 | .19 | .36 |
| Early Ocean | .16 | .16 | .20 | .20 |
| Ocean Natural | .50 | .50 | .50 | .50 |
| Ocean Harvest | 1.00 | 1.00 | .79 | .85 |
| In-river Harvest | .90 | .61 | .85 | .28 |
| Adult Passage | .70 | .91 | .55 | .83 |
| Pre-spawning | .85 | .85 | 1.00 | 1.00 |
| Lifetime Total (x10⁻³) | .46 | 2.69 | 3.51 | 3.56 |
| <i>Spawner/Spawner Ratio</i> | <i>1.16</i> | <i>6.74</i> | <i>6.04</i> | <i>6.13</i> |